

Prevalence and public health significance of anemia among non-pregnant, non-lactating tribal women of reproductive age (15-49 years) in India

Somnath Sau ^{1,2}, Dipak Kumar Tamili ³, Samiran Bisai ⁴, Koushik Das ⁵ and Balaram Das ^{1,*}

¹ Department of Physiology, Belda College, Belda-721424, Paschim Medinipur West Bengal, India.

² Department of Nutrition, Egra S.S.B. College, Egra-721429, Purba Medinipur, West Bengal, India.

³ Department of Zoology, Egra S.S.B. College, Egra-721429, Purba Medinipur, West Bengal, India.

⁴ Department of Anthropology & Tribal Studies, Public Health & Nutrition Research Unit, Sidho-Kanho-Birsha University, Purulia, West Bengal, India.

⁵ Department of Nutrition, Belda College, Belda-721424, Paschim Medinipur West Bengal, India.

Abstract

Background: In reproductive age, anemia is high burden and significant loss of maternal life in India. In general anemia largely a nutritional deficiency disorder, suffering about 50% of all women during reproduction.

Objectives: This review works elucidates the prevalence rate and highlighting the inadequacy or opportunities to refocus this area which are taken but lag behind these or could find other solutions.

Methods: The methodology of this review work are simply searching of published literature review, articles, periodicals, through online database, journal etc. The article selected from the year 2000 to 2020 for those work studied key areas of tribal non-pregnant, non-lactating reproductive women between 15-49 years age. This review study included final 20 eligible research papers those conducted on same age and physiological condition of tribal women.

Results: Out of 20 selected paper 7951 sample were gather and tabulated for analysis. The range of lowest 48 samples to 3923 big sample was varied. In general anemia prevalence calculated by 65.3% (95% CI: 64.3 – 66.4) in studied subjects.

Conclusion: Anemia is a worst situation among tribal women in India as per WHO classification of severity of anemia, indicating need based urgent steps.

Keywords: Anemia; Women; Micronutrient Deficiency; Iron deficiency; Tribal India

1. Introduction

The manifestation of a disease is much more of a worrying issue as the signs indicate underlying disturbed ease or literally what we call 'disease'. The appearance of ailment happens when indications developed so strong that the patient doings upon their indications [1]. Anemia is one such disease that is resulted for lower hemoglobin or hematocrit value. Red blood cells (RBC) containing a protein component, hemoglobin which transports oxygen to the whole body [2]. Lowering of Hb results the body may not get enough oxygen that they need to work the way they should. Clinically it can manifest as weakness or tiredness, unhappiness, and reduced mental function [3]. Anemia is a condition of reduced total

* Corresponding author: Balaram Das;

Department of Physiology, Belda College, Belda 721424, Paschim Medinipur West Bengal, India.

oxygen-carrying hemoglobin and/or red blood cells in blood [4][5]. According to WHO, below two standard deviations (SD) of Hb concentration or less from the population mean initiate anemia, which also vary with gender, age and populations [6].

Now a days, many developed countries like USA, fighting against different blood borne diseases like anemia. These countries will affect their population about 6% in many age groups from women to people who are risk for chronic health disorders [2]. WHO study group recommended hemoglobin level to diagnose anemia based on hemoglobin concentration in three stages- mild (110g/l to 119g/l), moderate (80 to 109g/l), and severe (< 80 g/l) among pre and post reproductive age of both sex, which are represented in (table 1) [7]. Women are more susceptible to anemia due to blood loss through menstruation and increase quantity of blood stock at pregnancy [2]. In the Global Nutrition Targets framework 2025, anemia is being key target, by achieving a 50% lessening of its occurrence in females with reproductive age endorsed by the World Health Assembly 2012 [8]. A systemic review study done in 2011 suggested that globally about 496 million (409–595 million) women are affected by anemia, which are non-pregnant and the worst affected regions were South Asia, West and Central Africa [9]. In the South-East Asia region, this number is approximately 190.6 million [9]. Demographic and Health Survey (DHS) report (2005-06) shows anemia (<12g/dl) prevalence in women and children in India is high among other South East Asian countries [10].

The most common type of anemia is iron-deficiency anemia across all ages [11]. Globally over 1.6 billion people affected by iron deficiency anemia most cases are from South Asia, Central and West Africa [9][10]. The most vulnerable people are indigenous groups of people around world. They represent approximately 476 million worldwide over more than 90 countries and make 6% of global population [12]. A significant maternal death is due to anemia and most cases due to low concentration of hemoglobin worldwide [9]. A systematic review data shows higher burden of anemia among indigenous people compare to general population [13]. Prevalence of nutritional anemia is largely distributed and affecting one quarter of world population [14]. In India, in view of anemia prevalence rate, there is large inequality of case distribution between inter-state and Union Territories. As per NFHS-2015-16 report, the percentages of about 80%, 76% and 66% women anemia cases respectively from Dadra and Nagar Haveli, Chandigarh, Andaman and Nicobar Island [15] and more than 60% anemia cases from Jharkhand, Haryana, West Bengal, Bihar and Andhra Pradesh [16]. The anemia prevalence of high-altitude states are Mizoram (25%), Manipur (26%), and Nagaland (28%). India is a large populous country and a committed nation to achieve Sustainable Development Goals (SDG) to reduce anemia prevalence at a set target level. Therefore, country needs more focus on women's anemia concern, precisely strata or sub-population level, where tribal women's reproductive age (15-49 years) is one among high priority groups [16].

As per census data 2011, India holds the estimated 10.4 crores tribal population which is roughly 8.6% of total population. They are inhabited mainly in forest and hilly regions of the country. Indian constitution has classified 705 Scheduled tribes and marked 75 as Particularly Vulnerable Tribal Groups (PVTGs), and that groups reside in 18 States and one Union Territory (UT) of India. The Scheduled Tribe male and female population is 52547215 and 51998501 respectively [16]. There is a need for more attention over both sexes' health concerns, especially for particularly vulnerable tribal groups.

India is a heavily populated country after china, although the per capita health expenditure cannot affordable. There is a health gap among different strata of the population; anemia is one priority health disorder. Approximately 50% of India's population is affected by anemia and women are more affected than men. In India, 20– 40 % maternal death is due to anemia [17]. Tribal women (15-49) are more ubiquitous in anemia cases, as NFHS-4 data shows prevalence rate is 59.9% [18]. There is a lack of broader coverage about anemia monitoring among Indigenous people worldwide. India has faced the same challenges [13]. Moreover, females are greater risk of developing anemia across different age groups and geographic regions worldwide [19]. In India, anemia is persistent, and the prevalence, not well drops, which are 53% of women (15-49years), is affected as per NFHS-4 survey report [18]. An article indicates anemia prevalence in women falls with an increased mean year of schooling and household income [14]. There is lack of sufficient published review literature among tribal communities in respect to proportion of population and number of vulnerable groups [20]. Since there is no wide scale coverage of published literature on anemia burden in tribal non-pregnant, non-lactating women age group 15- 49 years in India. Therefore, this contemporary review work is being studied for evaluation of the magnitude of the anemia burden and the community health consequence of tribal non-Pregnant non-lactating women (NPNL) throughout India.

2. Methodology

This review work focused on the overall anemia prevalence rate particularly tribal reproductive age (15 – 49years) women (non-Pregnant non- lactating women) in India. This review's methodology identifies relevant work on published papers on this subject in India's various states. The data on published works of literature is searched by electronic

platforms like Google, Google scholar, Pub Med, Researchgate, Springer from 2000 to 2020. Initially, the search items were nonspecific and huge on anemia but subsequently screen according to study goals. The keywords used to search relevant published articles include occurrence, determinants and distribution of anemia and its prevalence, tribal, micronutrients, WHO and UNICEF, India NFHS, state-wise tribal, particularly vulnerable tribal groups (PVTGs).

Table 1 WHO recommendation on Hemoglobin cut-off for anemia at sea level (g/dl)

Population	Non-anaemia	Mild	Moderate	Severe
6 - 59 months Children	≥11.0	10.0-10.9	7.0-9.9	< 7.0
5 - 11 years Children	≥11.5	11.0-11.4	8-10.9	< 8.0
12 - 14 years Children	≥12.0	11.0-11.9	8-10.9	< 8.0
15 years & above non-pregnant women	≥12.0	11.0-11.9	8-10.9	< 7.
Pregnant women	≥11.0	10.0-10.9	7-9.9	< 7.0
15 years & above Men		11.0-12.9	8-10.9	< 8.0

Sources: WHO VMNIS, 2011[21]

Table 2 WHO Classification of Public Health Significance of anemia

Category of public health significance	Prevalence of Anemia (%)
Severe	≥40.0
Moderate	20.0-39.9
Mild	5.0-19.9
Normal	≤4.9

Sources: WHO VMNIS, 2011[21]

Table 3 Age-sex specific RDA of Iron, Folic acid& Ascorbic acid (ICMR, 2020) [22]

Individuals	Age group	Body weight(kg)	Iron (mg/day)	Folate (μ/day)	Vit-C (mg/day)
Man		65	19	300	80
Women	NPNL	55	29	220	65
	Pregnant	55+10	40	570	80
	Lactation (0-6m, 7-12m)	-	23	330	115
		-		330	
Infants	0-6 m	5.8	-	25	20
	7-12m	8.5	3	85	27
Children	1-3yrs	11.7	8	110	27
	4-6yrs	18.3	11	135	32
	7-9yrs	25.3	15	170	43
Boys	10-12yrs	34.9	16	220	54
Girls	10-12yrs	36.4	28	225	52
Boys	13-15yrs	50.5	22	285	72
Girls	13-15yrs	49.6	30	245	66
Boys	16-18yrs	64.4	26	340	82
Girls	16-18yrs	55.7	32	270	68

Most of study article had been calculated based on WHO reference cut-off level of hemoglobin for drawing inferences about prevalence rate and degree of severity. However, some review articles were used NFHS reference cut-off point. To the knowledge of anaemia diagnosis, the WHO recommended Hb cut-off value at various age groups and grade of anemia i.e., mild, moderate or severe cases will be represented at Table 1&2[21]. Age and sex specific RDA of iron, folic acid and Vitamin-C are presented in table 3 [22].

3. Results

The total sample size of 20 selected articles sums into 7951 cases and the size of magnitude vary from minimum 48 to highest 3923cases (table 4) [23-42]. And within total sample the average percentage of prevalence is 65% at 95% confidence interval ranged lowest 64.3% to upper 66.4% anemic cases. The minimum range of anemia was 44.9% as reported by De et al, 2006 [24] from west Bengal (table 4). In contrary, highest rate of anemia (100%) was found among Munda women of west Bengal [23]. According to WHO reference criteria of anemia for public health significance, tribal women in India, are being severe affected with very high prevalence, indicating worst situation. Therefore, there is urgent need a nutrition education programme to improve their hemoglobin status.

Table 4 Summary of data collected from various studies on anemia

	Studied states	Ethnic groups	Year of data collection	Age groups (Years)	Sample Hb cut off	Methods of Hb. Estimation	Sample	Prevalence of anemia (%)
Ghosh&Bharati, 2003[23]	West Bengal	ST	2000	15-42	<12g/dl (WHO)	Sali's method	105	100
De et al., 2006[24]	West Bengal	ST	-	15 - 49	11g/dl (WHO)	Cell count	463	44.9
Unisa et al.,2010[25]	West Bengal	ST	2010	15 - 49	<12g/dl (NFHS-2)	Cyano-haemoglobin	137	66.0
De et al., 2011[26]	Assam	ST	-	15 - 45	11g/dl (WHO)	Cell count	173	76.3
Sreelakshmy et al.,2012[27]	Kerala	ST	2010	15 - 45	<12g/dl (WHO)	Cyano-haemoglobin	346	78.3
Kamath et al., 2013[28]	Karnataka	ST	2012	15 - 49	<12g/dl (NFHS-3)	HaemoCue	170	55.9
Shrinivasa, et al., 2014[29]	Kerala	ST	2013	15 - 45	<12g/dl (WHO)	Cyano-haemoglobin	347	96.5
Manna &Ghosh, 2014[30]	West Bengal	ST	2009	18 - 49	-	Cyano-haemoglobin	393	82.5
Bepari et al.,2015[31]	West Bengal	Lodha	-	18 - 45	<12g/dl	Haemo Cue	120	92.0
Nayak et al.,2016[32]	AndhraPradesh	ST	2014	15 - 49	<12g/dl	Cyano-hemoglobin	225	60.9
Ismail et al., 2016[33]	Kerala	ST	2014-15	≥ 18	<12g/dl (WHO)	Cyano-hemoglobin	113	64.6
Dabral et al.,2016[34]	Uttarakhand	Buksa	2014	15 - 45	<12g/dl	Sahli's method	112	64.3
Monsang& Sing, 2018[35]	Meghalaya	Garokhasi	-	15 - 49	<12g/dl (WHO)	HemoCue	150	92.0
Mahajan et. Al. 2019[36]	Gujarat	Kukna	2015-16	14 - 18	<12g/dl (WHO)	Drabkin's hemoglobin	168	45.3

Singh, 2019[37]	Meghalaya	ST	-	15 - 49	<12g/dl (WHO)	Sahli's method	103	89.3
Rohisha et al. 2019[38]	Kerala	ST	-	15 - 45	<12g/dl (WHO)	HemoCue	445	89.0
Chowdhury& Roy 2019[39]	West Bangal	Oraon	-	18-71	<12g/dl (WHO)	HemoCue	309	80.3
Dhanuka et al. 2019[40]	West Bengal	ST	2015-16	20-29	<12g/dl (WHO)	Cell count	48	62.5
Mahajan et al. 2019[41]	Jharkhand	Santal	2014-15	18-60	<12g/dl (WHO)	Cyano-hemoglobin	101	89.3
Rokade et al. 2020[42]	Maharastra	ST	2015-16	15-49	<12g/dl (NFHS4)	HemoCue	3923	54.0
Total							7951	65.3

4. Discussion

India has long-run policies and programs to curb anemia prevalence among all priority groups; despite that, it is still a country's burden [43]. The recent NFHS-5 data indicates anemia prevalence has not improved compared to NFHS-4 data across all categories [44]. Lack of anemia boosts the country's economic growth [45] and decreases faster as income increases [46].

Health for All's goal cannot be achieved fully unless given equal importance on health, especially tribal women's health and nutritional status [32]. Nutritional anemia is most common and easily avoidable [40]. Therefore, intake of micronutrient rich adequate diet is one way of the preventive method, mostly low-cost locally available diet for tribal families. In India, many studies are conducted to reveal the association between prevalence of anemia (%) and socioeconomic, environmental causes, dietary factors, poverty, illiteracy, low birth spacing, poor hygiene, and sanitation among tribal (NP-NL) women at 15 to 49 years of age [25][42].

Anemia is a significant community health problem between women age 15 – 49 during reproductive period (non-pregnant and pregnant) [47]. One-third of anemia is estimated in all women at reproductive age period (15-49 years). Globally the prevalence of anemia was 32.8 % (28.3% – 38.0%) among women in the reproductive age group and 32.5 % among women in non-pregnant on 2016. In South East Asia region, it was prevalent 45.8 % (39.1 – 51.2) in reproductive age women and 45.6% in non-pregnant women [48]. In India, according to 2016 data, anemia was 51.5% (41.48 – 58.75) and 51.43% (41.75- 58.55%) respectively among non-pregnant and reproductive age group women [44][49]. Anemia case is extensive among tribal women reproductive age group (15 – 49 years) [28]. The annual tribal health report in 2013- 14 suggested that anemia is prevalent in 65% tribal women in their reproductive age group (15-45 years)[49]. National Family Health Survey (NFHS) data from 2005 – 2020 on anemia (<12g/dl) suggests among non-pregnant women (15 – 49 yrs.) was 63.2% (2005-06), 62.8% (2015-16) and 71.7% (2019-20) respectively. Similarly, all women age 15 – 49 yrs. who are anemic was 63.2 % (2005-06), 62.5% (2015-16), and 71.4% (2019-20) respectively [44].

In 2011, WHO global report shows 48% (95% CI, 29 - 63) prevalence of anemia among Indian non-pregnant females (15-49years) at<12g/dl hemoglobin cut off level and level of public health significance of 2.5 at 8g/dl hemoglobin level (95% CI, 0.8 – 5.4) has been categorized as severe form. At the Global level, anemia prevalence was 29 % (95% CI, 23.9 – 34.8) in 2011[50]. According to NFHS 4 survey, the total sample size was 62695 tribal women, and the anemia prevalence rate was 59.9 % at <12g/dl hemoglobin level [18].

Health education and nutrition awareness are cost-effective strategies to reduce anemia prevalence [51]. Anemia is a preventable disease through dietary intervention, socioeconomic improvements etc. Dietary approach can become a successful tool in order to achieve healthy blood hemoglobin level. Vitamin-C is an important water-soluble substance, which increase non heam iron absorption by 20- 25%. It is a cost benefit dietary intervention [52]. However, a vitamin-C rich food is seasonal availability and some foods are expensive. In many tribal household a whole day's food supplies is cooked only once, providing two meals as per as 12 hours apart. Under such circumstance, it is difficult to meet ascorbic acid, iron and folic acid requirement unless a good source of these nutrients is added. Incorporation of iron

rich foods in the daily diet is the easiest and most necessary step. List of common Iron, Folic acid and vitamin-C rich foods are presented in table-5, 6, and 7 respectively [53].

Table 5 List of common iron rich foods (ICMR, 2017) [53]

SL No.	Iron rich foods	Scientific name	Amount (mg/100g edible portion)
1	Raisin dried, golden	<i>Vitisvinifera</i>	4.26±0.6
2	Raisin dried black	<i>Vitisvinifera</i>	6.81±0.91
3	Dates dry, pale brown	<i>Phoenix dactylifera</i>	3.20±0.45
4	Dates dry, dark brown	<i>Phoenix dactylifera</i>	4.79
5	Tamarind, pulp	<i>Tamarindusindicus</i>	9.16±1.71
6	Coriander leaves	<i>Coriandrum sativum</i>	5.30±1.55
7	Curry leaves	<i>Murraya koenigii</i>	8.67±0.09
8	Mint leaves	<i>Mentha spicata</i>	8.56±3.21
9	Asafoetida	<i>Ferula assafoetida</i>	15.68±4.51
10	Cardamom green	<i>Elettaria cardamomum</i>	8.33±1.44
11	Chillies red	<i>Capsicum annum</i>	6.23±0.79
12	Cloves	<i>Syzygium aromaticum</i>	9.41±2.10
13	Coriander seeds	<i>Coriandrum sativum</i>	17.64±6.74
14	Cumin seed	<i>Cuminum cyminum</i>	20.58±4.24
15	Pepper, black	<i>Piper nigrum</i>	11.91±3.48
16	Turmeric powder	<i>Curcuma domestica</i>	46.08±1.83
17	Gingelly seeds black	<i>Sesamum indicum</i>	13.9±1.60
18	Mustard seeds	<i>Brassica juncea</i>	13.49±3.95
19	Egg, poultry yolk boiled	<i>Gallus gallus</i>	4.92±0.33
20	Poultry chicken liver	<i>Gallus gallus</i>	9.92
21	Goat spleen	<i>Capra aegagrus</i>	51.41±17.21
22	Beef spleen	<i>Bos taurus</i>	31.68±4.50
23	Pork spleen	<i>Sus scrofa</i>	27.21±6.43
24	Pork liver	<i>Sus scrofa</i>	20.74±7.24
25	Amaranth seed black	<i>Amaranthus cruentus</i>	9.33
26	Ragi	<i>Eleusine coracana</i>	4.62±0.36
27	Rice flakes	<i>Oryza sativa</i>	4.46±0.81
28	Puffed rice	<i>Oryza sativa</i>	4.55±1.03
29	Wheat flour atta	<i>Triticum aestivum</i>	4.10±0.67
30	Bengal gram, dal	<i>Cicer arietinum</i>	6.08±0.27
31	Bengal gram whole	<i>Cicer arietinum</i>	6.78±0.75
32	Horse gram Whole	<i>Dolichos biflorus</i>	8.76±1.16
33	Lentil whole brown	<i>Lens culinaris</i>	7.57±0.67
34	Soyabean brown	<i>Glycine max</i>	8.29±0.51

35	Beet green	<i>Beta vulgaris</i>	5.8±0.57
36	Fenugreek leaves	<i>Trigonellafoenumgraecum</i>	5.69±1.37
37	Drumstick leaves	<i>Moringaoleifera</i>	4.56±1.09
38	Pumpkin leaves	<i>Cucurbita maxima</i>	5.58±0.39
39	Onion stalk	<i>Allium cepa</i>	3.09±0.54

Table 6 List of common total folate rich foods (ICMR, 2017) [53]

Sl. No.	Total folates (B9) rich foods	Scientific name	Amount (μ/100g edible portion)
1	Bengal gram whole	<i>Cicerarietinum</i>	233±12.9
2	Black gram whole	<i>Phaseolusmungo</i>	134±14.2
3	Cow pea brown	<i>Vignacatjang</i>	231±27.3
4	Cow pea white	<i>Vignacatjang</i>	249
5	Field bean white	<i>Phaseolus vulgaris</i>	289±27
6	Moth bean	<i>Vignaconitifolia</i>	349±10.8
7	Rajmah brown	<i>Phaseolus vulgaris</i>	330±29.6
8	Red gram whole	<i>Cajanuscajan</i>	229±19
9	Soyabean brown	<i>Glycine max</i>	297±26.1
10	Parsley	<i>Petroselinumcrispum</i>	197±13.9
11	Drumstick leaves	<i>Moringaoleifera</i>	42.89±5.31
12	Mustard leaves	<i>Brassica juncea</i>	110±6.6
13	Spinach	<i>Spinach oleracea</i>	142±10.3
14	Tamarind leaves tender	<i>Tamarindusindica</i>	91.82±9.56
15	Capsicum, green	<i>Capsicum annuum</i>	51.85±3.38
16	Capsicum red	<i>Capsicum annuum</i>	62.54±2.15
17	Ladiesfinger	<i>Abelmoschusesculentus</i>	63.68±10.76
18	Mango ripe himsagar	<i>Magniferaindica</i>	90.98±6.12
19	Papaya ripe	<i>Carcia papaya</i>	60.90±6.64
20	Beetroot	<i>Beta vulgaris</i>	97.37±7.06
21	Curry leaves	<i>Murrayakoenigii</i>	117±19.3
22	Garlic, big clove	<i>Allium sativum</i>	85.77±15.61
23	Mint leaves	<i>Menthaspicata</i>	106±6.3
24	Poppy seeds	<i>Papaversomniferum</i>	78.73±7.90
25	Gingelly seeds brown	<i>Sesamumindicum</i>	92.63±5.90
26	Paneer	----	93.31±14.37
27	Khoa	-----	94.25±8.57
28	Egg yolk raw poultry	<i>Gallus gallus</i>	112±6.1
29	Egg yolk boiled poultry	<i>Gallus gallus</i>	110±6.1
30	Sheep liver	-----	206±26.8
31	Beef liver	<i>Bostaurus</i>	1744±71.2

32	Calf liver	-----	1473
33	Aluva	<i>Parastromateusniger</i>	1132±159
34	Betki	<i>Latescalcarifer</i>	2079
35	Bombay duck	<i>Harpadonnehereus</i>	2784
36	Hilsa	<i>Tenualosailisha</i>	2875
37	Silver carp	<i>Hypophthalmichthysmolitrix</i>	2462
38	Carb	<i>Menippemercenaria</i>	2304
39	Octopus	<i>Octopus vulgaris</i>	2087
40	Catla	<i>Catlacatla</i>	1926±277
43	Rohu	<i>Labeorohita</i>	1263±101
44	Tiger prawns	<i>Macrobrachium sp.</i>	1875

Table 7 List of common vitamin-C rich foods (ICMR, 2017) [53].

Sl. No.	Vitamin-C rich foods	Scientific name	Amount (mg/100g edible portion)
1	Agathi leaves	<i>Sesbaniagrandiflora</i>	121
2	Amaranth leaves, red	<i>Amaranth gangeticus</i>	86.20
3	Amaranth leaves	<i>Amaranth viridis</i>	179
4	Brussels sprouts	<i>Brassica oleraceavar.gemmifera</i>	89.45
5	Drumstick leaves	<i>Moringaoleifera</i>	108±16.7
6	Parsley	<i>Petroselinumcrispum</i>	133±16.3
7	Ponnaganni	<i>Alternantherasessilis</i>	103
8	Raddish leaves	<i>Raphanussativus</i>	65.76±18.69
9	Bitter gourd, jagged, smooth ridge elongate	<i>MomordicaCharantia</i>	54.30
10	Capsicum green	<i>Capsicum Annuum</i>	123±7.8
11	Capsicum red	<i>Capsicum Annuum</i>	112±5.5
12	Capsicum yellow	<i>Capsicum Annuum</i>	127±12.5
13	Drumstick	<i>Moringaoleifera</i>	71.86±19.13
14	Knol-Khol	<i>Brassica oleracea</i>	64.70±10.78
15	Mango green raw	<i>MagniferaIndica</i>	
16	Currants, black	<i>Ribesnigrum</i>	182
17	Gooseberry (Amla)	<i>Emblicaofficinalis</i>	252±30.4
18	Guava, white flesh	<i>Psidiumguajava</i>	214±13.6
19	Guava, pink flesh	<i>Psidiumguajava</i>	222±27
20	Manila tamarind	<i>Pithecellobiumdulce</i>	55.78
21	Strawberry	<i>Fragaria X ananassa</i>	50.20±4.97
22	Cillies, green- all varieties	<i>Capsicum annum</i>	94.07±11.67

Anemia is a major public health problem in India [40]. About 65% of tribal females having age between 15-49 years are anemic. A significant proportion of anemia is result of nutritional inadequacy. Therefore, anemia is preventable disease. NFHS technical report 2009 had shown average daily household intake of food stuffs by ST population does not meet RDA (%) such as green leafy vegetables 56%, other vegetables 69%, milk and milk products 14%, fats and oils 50%, sugar and jaggery 30% and pulses 75% of RDA. Average daily household nutrient intake by ST population was below than RDA (%), such nutrients are iron 44%, folic acid 51%, 36% riboflavin, 46% vitamin A and protein 78% of RDA as per tribal health report 2018, India [49].

5. Conclusion

From this review it is evident that, a high burden of anemia prevalence among tribal NPPL (15-49yrs.) reproductive women in India (Table-4), since 2000- 2016. So, it creates matters of concern and a severe public health thread in future. Dietary intervention, good personal hygiene and sanitation have potential impact on anemia reduction as evidence by several studies of this review. In India IFA supplementation (iron + initiatives 2013) is fail to reduce the burden of IDA at implementation level among target groups [54]. As per WHO for India, health services enhancement and food supplementation are effective strategy for anemia prevention. With that Community based screening for Hb level, monitoring of IFA distribution especially adolescent & reproductive women at ground level will produce better results. Above all these efforts along with community participation, Individual felt needs and strong political commitment will necessary to assure desire outcome very soon. Recommendation for locally available low-cost seasonal foods preferentially vitamin C, iron, folic acid rich food items etc. are guiding principles for better anemia management. This review may be best utilized as an educational tools and guidance's by community health workers, nutritionists, nurses and policy makers.

Compliance with ethical standards

Acknowledgments

The authors express gratefulness to the Belda College, Belda, Paschim Medinipur for providing the facilities to execute the review study. We are heartily thankful to Dr. Sandeep Kumar Dash, Department of Physiology, University of Gourbanga, Malda, India.

Disclosure of conflict of interest

Authors declare that there are no conflicts of interests.

References

- [1] Prince AE, Berkman BE. When does an illness begin: genetic discrimination and disease manifestation? The Journal of law, medicine & ethics: a journal of the American Society of Law, Medicine & Ethics. 2012; 40(3): 655–664.
- [2] Felson S. anaemia WebMD, [Internet source] 2020, August 11, available from: <https://www.webmd.com>.
- [3] Smith RE Jr. The clinical and economic burden of anemia. American Journal of Managed Care. 2010; 16(3):59-66.
- [4] Urs Giger PD, Dr. Med. Vet., M.S., F.V.H., DACVIM, DECVIM, anaemia. In: Deborah C. Silverstein, Kate Hopper. editor. In Small Animal Critical Care Medicine, W.B. Saunders; 2009: 518-523
- [5] Rodak BF. Hematology: clinical principles and applications (3rd ed.). Philadelphia: Saunders; 2007: p. 220. ISBN 978-1-4160-3006-5. Archived from the original on 2016-04-25.
- [6] De Benoist B, Cogswell M, Egli I, McLean E. Worldwide prevalence of anaemia 1993-2005. WHO Global Database of Anaemia; 2008:1–40.
- [7] World Health Organization. Hemoglobin concentrations for the diagnosis of anemia and assessment of severity; 2011: p-6, Accessed August 4, 2017 available from <http://www.who.int/vmnis/indicators/haemoglobin.pdf>.
- [8] World Health Organization. Intervention by global target. eLENA. Department of Nutrition for Health and Development; 2012-13
- [9] Stevens. GA, Finucane. MM, De-Regil LM, Paciorek CJ, Flaxman. SR, Branca. F, Peña-Rosas. JP, Bhutta. ZA, Ezzati. M. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia

in children and pregnant and non-pregnant women for 1995–2011: a systematic analysis of population-representative data. *The Lancet Global Health*. 2013; 1(1):16-25.

- [10] Kothari. MT and Abderrahim. N. Nutrition Update 2010. DHS Nutritional Reports No. 4. United States Agency for International Development. Calverton, Maryland, USA: ICF Macro. 2010. Available at <http://dhsprogram.com/pubs/pdf/NUT4/NUT4.pdf>.
- [11] Johnny Amer. A Retrospective Study Using Mentzer Index for Prevalence of Iron Deficiency Anemia among Infants Visiting Maternal Centers at the Age of One Year, *Anaemia*, 2022; Vol. 2022: p-5, Article ID 7236317, <https://doi.org/10.1155/2022/7236317>.
- [12] United Nations, In: International Day of the World's Indigenous Peoples. 2021. <https://www.un.org/en/observances/indigenous-day>.
- [13] Khamalia AZ, AimoneAM, Zlotkin SH. Burden of anemia among indigenous populations. *Nutrition Reviews*. 2011; 69(12):693-719.
- [14] Leite. M, Cardoso. A., Coimbra. C., Welch. J., Gugelmin. S., Lira. P., Horta. B., Santos. R. and Escobar. A. Prevalence of anemia and associated factors among indigenous children in Brazil: results from the First National Survey of Indigenous People's Health and Nutrition. *Nutrition Journal*. 2013; 12:69
- [15] Adil Akhzer, 2015-16 National Family Health Survey Report Anaemia among kids aged 6-59 months highest in Chandigarh: Report; *The Indian Express*; January 12, 2018.
- [16] Census of India, Office of the Registrar General and Census Operation, Ministry of Home Affairs, Government of India, New Delhi, India, 2011.
- [17] Jadhav BN. A critical review on iron deficiency anemia in female population of developing India. *International Journal of Fauna and Biological Studies*. 2016; 3:116-119.
- [18] International Institute for Population Sciences (IIPS) and ICF. National Family Health Survey (NFHS-4), 2015-16: India. Mumbai: IIPS. 2017.
- [19] Kassebaum NJ, Jasrasaria R, Naghavi M, Johns N, Lozano R, Regan M, Weatherall D, Chou DP, Eisele TP, Flaxman SR, Pullan RL, Brooker SJ and Murray CJL. A systematic analysis of global anemia burden from 1990 to 2010. *Blood*. 2014; 123(5):615–624.
- [20] Rakesh P. S. Prevalence of Anaemia in Kerala State, Southern India - A Systematic Review. *Journal of clinical and diagnostic research*. 2017; 11(5):1–4.
- [21] WHO. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and Mineral Nutrition Information System. Geneva, World Health Organization, 2011 (WHO/NMH/NHD/MNM/11.1) (<http://www.who.int/vmnis/indicators/haemoglobin>).
- [22] ICMR- NIN Expert group on Nutrient Requirement for Indians, Recommended Dietary Allowances (RDA) and Estimated Average Requirements (EAR)-2020.
- [23] Ghosh R, Bharati P. Haemoglobin status of adult women of two ethnic groups living in a peri-urban area of Kolkata city, India: a micro-level study. *Asia Pac J ClinNutr*. 2003; 12(4):451–59.
- [24] Madhusnata De, Ajanta Halder, Sandeep Podder, Rinini Sen, Shila Chakrabarty, Bani Sengupta, Tulika Chakraborty, Urmisha Das & Geeta Talukder. Anemia and hemoglobinopathies in tribal population of Eastern and North-eastern India. *Hematology*. 2006; 11:5-6, 371-373, DOI: 10.1080/10245330600840180
- [25] Unisa S., Chattopadhyay A., Ganguly S., Roy T.K. Causes of Anemia among Women in the Eastern Region of India: A Study of West Bengal. Working Paper No. 3, International Institute for Population Sciences, Mumbai. 2010.
- [26] De M., Halder A., Chakraborty T., Das U., Paul S., De A., Banerjee J., Chatterjee T., De S. Incidence of anemia and effect of nutritional supplementation on women in rural and tribal populations of eastern and North-eastern India. *Hematology*. 2011; 16(3):190-192.
- [27] Sreelekshmy PR, Vijaykumar K, Anish TS, Srinivasa BM, Sheela's , Abraham A . Anaemia & BMI of non-pregnant tribal Women of Reproductive age group in palakkad District of Kerela. *Ind J of Pre Soc-Med*. 2012, 43 (1): 35-41.
- [28] Kamath R, Majeed JA, Chandrasekaran V, Pattanshetty SM. Prevalence of anemia among tribal women of reproductive age in Udupi Taluk, Karnataka. *Journal of Family Medicine and Primary Care*. 2013; 2(4):345-348.

- [29] Shrinivasa BM, Philip RR, Krishnapali VK, Suraj A, Sreelakshmi PR. Prevalence of anemia among tribal women of reproductive age-group in Wayanad district of Kerala. *International Journal of Health and Allied Sciences*. 2014; 3(2):120-124.
- [30] Manna. K. Prabir and Ghosh. Debidas. Prevalence of Anemia among Adolescent Girls and Adult Women of Tea Garden Areas of Darjeeling and Jalpaiguri Districts of West Bengal. *J Life Science*. 2014; 6(1, 2): 33-39.
- [31] Bepari M., Pal A., Maity P., Choudhury SM. Nutritional and health status of adult women of the Lodha tribal population of PaschimMidnapore, West Bengal, India: compared with nontribal women. *East African J Public Health*. 2015; 12(1):988-996.
- [32] Nayak MSDP and Sreegiri S. A study on nutritional status of tribal women in Visakhapatnam district, Andhra Pradesh, India. *International Journal of Community Medicine and Public Health*. 2016; 3(8):2049-2053.
- [33] Ismail IM., Kahkashan A., Antony A., Sobhith VK. Role of socio-demographic and cultural factors on anemia in a tribal population of North Kerala, India. *International Journal of Community Medicine and Public Health*. 2016; 3(5):1183-1188.
- [34] Dabral M. and Kothiyal P. Prevalence of anemia among reproductive age group tribal women in Uttarakhand, India. *Indian Journal of Pharmaceutical Science & Research*. 2016; 6(1):47-50.
- [35] Monsang SE and Sing N. Prevalence of Anaemia and Nutritional Knowledge among Tribal Women of Reproductive Age Group of Meghalaya, India. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7(10):1221-1229.
- [36] Mahajan N, Kshatriya GK. Trends of Nutritional Anaemia Among Adolescents of Kukna Tribal Community of Gujarat, India. *Online Jorنال of Health and Allied Sciences*. 2019; 18(2):1-6.
- [37] Singh. N. Nutritional Status of Tribal Women of Reproductive Age Group in Naugarh Block, Chandauli District, India. *International Journal of Current Microbiology and Applied Sciences*. 2019; 8(7): 953-966.
- [38] Rohisha IK, Jose TT, Chakrabarty J. Prevalence of anemia among tribal women. *Journal of Family Medicine and Primary Care*. 2019; 8(1):145-147.
- [39] Chowdhury TK. and Roy SK. Prevalence of anaemia and associated factors among Oraon females. *Anthropological Review*. 2019; 82(1):15-27.
- [40] Dhanuka A., Goswami BK., Goswami SB., Chakrabarti S. Profile of nutritional anemia and its correlation with serum iron, Vitamin B12, and folic acid level among the tribal population of northern districts of West Bengal, India. *Archives of Medicine and Health Sciences*. 2019; 7:201-205.
- [41] Mahajan N, Kumari A and Kshatriya GK. Prevalence of undernutrition and gender disparities among Kukna and Santal tribal population groups of India. *Research Journal of Humanities and Social Sciences*. 2019; 10(2):351-358.
- [42] Rokade S, Mog M and Mondal NA. Nutritional status among tribal women in Maharashtra, India: Spatial variations and determinants. *Clinical Epidemiology and Global Health*. 2020; 8:1360-1365.
- [43] Anand T., Rahi M., Sharma P., Ingle GK. Issues in prevention of iron deficiency anemia in India. *Nutrition*. 2014; 30(7-8):764-70
- [44] National Family Health Survey (NFHS – 5), (2019-21), http://rchiips.org/nfhs/factsheet_NFHS-5.shtml.
- [45] The World Bank national accounts data, and OECD National Accounts data files. GDP growth (annual %), 2018.
- [46] Alderman H. and Linnemayr S. Anemia in low-income countries is unlikely to be addressed by economic development without additional programs. *Food and Nutrition Bulletin*. 2009; 30:265-269
- [47] Takana M Silubonde, Cornelius M Smuts, Lisa J Ware, Glory Chidumwa , Linda Malan, Stephen Lye ,Shane A Norris. Determinants of Anaemia Among Women of Reproductive Age in South Africa: A Healthy Life Trajectories Initiative (HeLTI); 2021 <https://doi.org/10.21203/rs.3.rs-948470/v1>
- [48] WHO (n.d). Prevalence of anaemia in women of reproductive age (aged 15-49) (%). Retrieved April 10, 2022, from [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-anaemia-in-women-of-reproductive-age-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-anaemia-in-women-of-reproductive-age-(-)).
- [49] Tribal Health in India – Bridging the gap and a roadmap for the future. Report of the expert committee on tribal health. Ministry of Health and Family Welfare, Government of India, New Delhi; 2018.

- [50] World Health Organization. The Global prevalence of anemia in 2011. Geneva: World Health Organization; 2015: ISBN 978 92 4 156496 0 (NLM classification: WH 155).
- [51] Mehrotra M, Yadav S, Deshpande A, Mehrotra H. A study of the prevalence of anemia and associated sociodemographic factors in pregnant women in Port Blair, Andaman and Nicobar Islands. *Journal of Family Medicine and Primary Care*. 2018; 7(6):1288-1293.
- [52] De Maeyer E.M et.al, Prevention and controlling iron deficiency anaemia: the contribution of primary health care, a guide for health administrators and programmemanagers, A draft document prepared for the International Nutritional Anemia Consultative Group (INCAG), 17th Jul 1997. Washington, DC: USA: INCAG.Nutrition Foundation; Mimeo.1998.
- [53] Longvah T, Ananthan R, Bhaskarachary K and Venkaiah. Indian food composition table, Telangana (India): National institute of Nutrition; 2017 <http://www.ifct2017.com/frame.php?page=food>.
- [54] Rai et al. Iron-deficiency anaemia among women in India, *WHO South East Asia Journal of Public Health*, April 2018; 7(1): 18-2.