

Phenotypic traits of three local cassava (*Manihot esculenta* Crantz) genotypes cultivated by farmers in Rivers State, Nigeria

Josephine U Agogbua * and Barida Biranen

Department of Plant Science and Biotechnology, University of Port Harcourt, Nigeria.

Abstract

The study aimed to elucidate the morphological and anatomical traits for varietal delimitation and identification of three local cassava (*Manihot esculenta* var. aguregbu, var. cagbo and var. ekagon) varieties cultivated by farmers in Obio Akpor local government area of Rivers state. Twenty (20) qualitative and six (6) quantitative traits from the descriptors for cassava germplasm was used for morpho-characterization. The results obtained revealed phenotypic similarity for 15 qualitative traits among the three varieties. Trait differences was observed in the petiole color (red and green), prominence of foliar scars, and shape of stem, level of branching, root shape and constrictions. The distance between leaf scars was classed as short for all varieties but the metric value ranged from 1.6 cm for ekagon to 4.0 cm for aguregbu and cagbo. Other quantitative variation observed include petiole length (11.1 to 15.0 cm) and storage root length which was classified as medium (≤ 30 cm) and long (> 30 cm). The number of leaf lobes was seven for all varieties. The epidermal traits showed a uniseriate epidermis with hypostomatic distribution. The stem, petiole, and midrib anatomy was similar in terms of tissue type and arrangement. However, the number of vascular bundles in the petiole was more in aguregbu (10) than in cagbo (8) and ekagon (8). The variable traits identified in this study provide diagnostic features that could aid in the identification of the three cassava genotypes.

Keywords: *Manihot esculenta* var Aguregbu; Cagbo; Ekagon; Petiole; Phenotypic traits

1. Introduction

Manihot esculenta Crantz commonly called cassava is major staple in many developing countries of Africa, South and Central America, India and Southeast Asia [1]. It is known for its unique ability to grow and produce well in low fertility soil, withstand attack by locusts and other insect pests, and tolerate drought conditions. Cassava serves as an important staple food and can also be used directly or as protein source in the concentrate mixture in animal feed [2]. Naturally fermented cassava starch is used in bread-making [3]. In Nigeria and other African countries, tapioca and fufu are made from starchy cassava root flour while garri is made from fermentation of root [4]. The seeds contains 90% unsaturated oil which might possibly be used as a fuel for pre- combustion diesel engines [5]. The production and growth of cassava is severely limited in certain semi- arid areas such as north-eastern Brazil and Northern Nigeria, due to drought conditions [6]. The bitter variety of *Manihot* root is used to treat diarrhoea and malaria [7]. The leaves are used to treat hypertension, headache, and pain [8]. Cubans commonly use cassava to treat irritable bowel syndrome, the paste is eaten in excess during treatment [9]. It has been reported that cassava may have anti-cancer properties [7]. Genes isolated from the plant have been already used to eradicate brain tumours in laboratory rats [10].

Anatomical traits in foliar epidermis, stem and petiole which are highly conserved and not affected by the environment have been extensively utilized for identification of plants and elucidation of genetic relationships. Stem [11] and petiole [12] anatomy have been used for taxonomic delimitation of wild cassava genotypes, cultivars and hybrids. The objective

* Corresponding author: Josephine U Agogbua
Department of Plant Science and Biotechnology, University of Port Harcourt.

of this research was to determine the relationship of three local varieties of *Manihot esculenta* var. aguregbu, cagbo and ekagon cultivated by farmers in Rivers State, Nigeria using phenotypic markers.

2. Material and methods

2.1. Sample collection

The three local varieties of *Manihot esculenta* used in this study (*Manihot esculenta* var. aguregbu, *Manihot esculenta* var. cagbo, *Manihot esculenta* var. Ekagon) were collected from a farmers field in Rumuosi, in Rivers State, with GPS location of Latitude 04°55'14.8" North, longitude 006°55'07.7" East.

2.2. Morphological traits

Standard morphological and agronomic descriptors for characterization of cassava germplasm [13] was used to score vegetative traits in the field. Overall morphology of the plants were recorded by photography. The quantitative traits measured were petiole length, leaf width, leaf length, number of lobes, distance between leaf scars and storage root length.

2.3. Anatomical traits

Fresh specimen of leaves, midrib and stems were collected from each of the variety in the field and used for the anatomical studies.

2.3.1. Foliar Epidermis

The upper and lower epidermal layers of fresh matured leaves were peeled and stained with 1% safranin or alcian blue, rinsed with distilled water to remove excess stain and then mounted in a drop of pure glycerol on clean glass slides and viewed using a light microscope. Photomicrographs of both adaxial and abaxial surfaces were taken with the aid of a digital camera.

2.3.2. Stem, Petiole and midrib

Stem, petiole and midrib were collected and hand sectioned using the method of [14]. Sections were placed in a Petri dish that contained water and subsequently collected using a camel brush, and placed on a clean glass slide and then stained with 1% safranin or alcian blue and then rinsed with the distilled water. The sections were then mounted on slides with a drop of glycerin, viewed under the microscope and photographed.

3. Results

3.1. Morphological studies

The qualitative and quantitative descriptors scored for all morphological traits in the three varieties are on table 1 and figure 1.

3.1.1. *Manihot esculenta* var. Aguregbu

The plant type is cylindrical, it has a one level branching system, it has a dark green color of fully expanded leaf, the shape of the central lobe is lanceolate, it has a light green leaf vein color, the petiole color is totally pigmented with a pink color, the stem is dark brown, the storage root shape is conical-cylindrical, it has prominent leaf scars, storage root constrictions is present with a storage root length of more than 30 cm.

3.1.2. *Manihot esculenta* var. Ekagon

The plant type is open, it has a three branching level type, the first fully expanded leaf is light green in color, the leaf shape is lanceolate, the leaf vein color is light green, the petiole color is green without pigmentation, the stem color is dark brown, the root storage system is dark brown, it has a fusiform storage root shape, there is presence of root constrictions, it has little leaf scar prominence, the storage root length is long more than 30 cm.

3.1.3. *Manihot esculenta* var. Cagbo

The plant type is open, it has a one level branching system, the first fully expanded leaf is dark green, it has a lanceolate leaf shape, the petiole color is dark green, it has a dark brown stem color, there is absence of root constrictions, it has

moderately prominent leaf scars, the storage root length is medium with a length of 20-30 cm, the storage root shape is conical.

Table 1 Morphological characters of three local varieties of cassava evaluated in the farmer's field

Variety			
DESCRIPTORS	Aguregbu	Cagbo	Ekagon
Qualitative Traits			
Plant shape	Open	Cylindrical	Open
Color of apical leaf	Light green	Light green	Light green
Pubescence on apical leaf	Absent	Absent	Absent
Shape of central leaflet	Lanceolate	Lanceolate	Lanceolate
Leaf color	Dark green	Light green	Dark green
Petiole color	Red	Green	Green
Lobe margin	Smooth	Smooth	Smooth
Color of leaf vein	Green	Green	Green
Orientation of petioles	Horizontal	Horizontal	Horizontal
Prominence of foliar scars	Prominent	Semi prominent	Semi prominent
Color of stem exterior	Dark brown	Dark brown	Dark brown
Color of stem cortex	Dark green	Light green	Light green
Color of stem epidermis	Dark brown	Light brown	Light brown
Growth habit of stem	Straight	Straight	Straight
Color of end branches	Green-purple	Green	Green
Shape of stem	Cylindrical	Open	Open
Branching habit	Dichotomous	Dichotomous	Dichotomous
Level of branching	One	One	Three
Root constrictions	Present	Absent	Present
Root shape	Conical-cylindrical	Conical	Fusiform
Quantitative Traits			
No. of leaf lobes	7	7	7
Leaf length (cm)	12.9	11.1	14.1
Leaf width (cm)	18.5	20.7	23.4
Petiole length (cm)	11.1	14.9	15
Distance between Leaf scars(cm)	4.0 Short	3.9 Short	1.6 Short
Storage root length (cm)	>30 long	≤30 medium	>30 long

Table 2 Epidermal traits of the upper and lower leaf surface of three local cassava varieties

Variety						
DESCRIPTORS	Aguregbu		Cagbo		Ekagon	
	Upper	Lower	Upper	Lower	Upper	Lower
Foliar Epidermal Traits						
Shape of epidermal cells	Regular	Irregular	Regular	Irregular	Regular	Irregular
Stomata distribution	Hypostomatic		Hypostomatic		Hypostomatic	
Trichomes	Absent	Absent	Absent	Absent	Absent	Absent




Figure 1 Habit of three local cassava varieties- aguregbu, cagbo and ekagon

3.2. Anatomical Studies

Epidermal anatomy

This is divided into two the adaxial which is the upper part of the leaf, and abaxial which is the lower part of the leaf.

- Leaf-Adaxial

The result revealed a glabrous uniseriate adaxial epidermis with regular shaped cells for the three varieties. No stomata was observed in the upper epidermis (Figure 2).

- The Leaf-Abaxial

The adaxial epidermis revealed a glabrous uniseriate epidermis with irregular shaped cells. The stomata is mostly anomocytic with a stomatal index of 13.02. The guard cells are elliptical with mostly 5–6 subsidiary cells (Figure 2)..

Stem anatomy

The stem is pentagonal in cross or transverse section, having 5 ridges and 5 furrows. There is presence of epidermal layer, with the absence of trichomes. Endodermis is found to enclose the vascular bundles. The vasculature of the stem shows the arrangement of the vascular bundles with respect to number of furrows and ridges present. The vascular bundles are collateral having phloem and xylem linked by the cambium. Pith is distinct with a well-developed parenchymatous cells. These cells interspersed with air spaces to form a characteristic pattern of pith tissues. The stem cross section is seen in figure 3.

Petiole Anatomy

The outline of the petiole was observed to be circular in shape. The epidermal cells are uniseriate, with the absence of trichomes. Endodermis is found to surround the vasculatures in which the vascular bundles are arranged. The number of vasculatures present varies among the varieties and this variation ranges from 8 to 10. The arrangement of vascular

bundle is circular in nature. The vascular bundles vary in shape and are collateral, having phloem and xylem linked by cambium. Pith is also present and consists of well-developed parenchymatous cells which are irregularly arranged. The number of vascular bundles in the petiole of aguregbu was more (10 vascular bundles) than var. cagbo and var. ekagon which had 8 vascular bundles (Figure 3). Trichomes were not observed in any of the varieties.

Midrib anatomy




Figure 2 Foliar epidermis of aguregbu, cagbo and ekagon cassava varieties




Figure 3 Transverse section of Stem, Petiole and Midrib of aguregbu, cagbo and ekagon cassava varieties

The midrib is highly conspicuous and U-shape in cross section with angular collenchyma cells on both upper and lower surfaces. The shape varies among the varieties. The vascular bundles are collateral and are arranged in U shape. The

bundle sheaths are mainly parenchymatous and are also arranged in u shape. The parenchyma cells filled the space between the collenchyma and the vascular bundles. The vascular bundles in the midrib is collateral with the phloem towards the lower leaf surface. Trichomes were absent (Figure 3).

4. Discussion

This present study provides information on the morphological and anatomical studies of three varieties of cassava. The results obtained from the morphological studies, showed that the variety aguregbu exhibited the highest leaf number and are also branched. This indicates that the variety will exhibit higher tendency of carrying out photosynthetic activities, which may also lead to an increase in yield (15).

The morphology of the cassava varieties provided diagnostic features (16). Some of the varieties like aguregbu consists of petiole that are highly pigmented (17) while varieties like ekagon and cagbo consist of petioles that are not pigmented but are greenish in color. This variation in petiole pigmentation distinguishes Aguregbu from the other two varieties. Most *Manihot* species have long petioles (18). The petioles and tissues contain the same arrangement of those present in the stem of *Manihot* (19). The petiole of dicotyledons often have similar provision stem (20). Trichomes were not observed in this study. The presence of trichomes, thick cuticles, crystals and anthocyanins help to minimize water loss in dry environment [21]. The existence of trichomes is related to the process of speciation that favor the trichomes in low humidity environment (22). The presence of anthocyanins leads to photoinhibition and can accumulate on the stem of the plant in response to stress induced by cold or heat [23].

Morphological and Anatomical features, together with other sources of evidence such as pigmentation, stem leaf scars, branching system and stem root constrictions, shows that the varieties of cassava (*Manihot esculenta* Crantz) worked on, are mesophytes and upright [24], with highly evolving characteristics of a dicot. The hypodermis plays an important role in drought resistance, and probably functions as a water storage tissue in such species [19]. These anatomical changes may occur in plants as an adaptation to drought stress[25].

All the varieties studied, expressed an identical stem color [26]. Results obtained from the anatomical studies on the three varieties, showed a useful characteristic with respect to the presence of branching system and arrangement of the vascular bundles in both petiole and midrib [27]. In the stem anatomy, the anatomical evidence reveals the relatedness of these. Some specific variations were observed in the midrib and petiole. These variations are of great taxonomic importance [28].

In petiole anatomy, variety aguregbu expressed the highest number of vascularbundles (10) and are branched, followed by Cagbo and Ekagon which both had 8 vascular bundles each. A continuous layer of collenchyma was observed (which is involved in mechanical support) of variable thickness in the cortex below the chlrenchyma. Pericyclic fibers in the cortex forms two to three continuous layers [29]. Plants under drought stress, the fibers have the same distribution, and the number of layers increases. Sclerenchyma helps to prevents water loss, which explains the adaptability of species in tropical regions, such as *Manihot*, to drought stress [19]. Laticifers were observed in the cortex, phloematic tissue, xylem, and on the periphery of the pith, and is a common characteristic of the Euphorbiaceae.

5. Conclusion

In this study, the distinguishing traits among the three varieties were petiole color, petiole length, level of branching, storage root constrictions and number of vascular bundles in petiole. These variable traits identified in this study provide diagnostic features that could aid in the identification of the three cassava genotypes.

Compliance with ethical standards

Acknowledgments

The authors acknowledge the local farmers for permission to use their fields and providing the plant materials used for this study.

Disclosure of conflict of interest

The authors declare no conflict of interest.

References

- [1] Temesgen M, Retta N. Nutritional potential, health and food security benefits of taro *Colocasia esculenta* (L.): A review. *Food Science and Quality Management*. 2015; 36:23-30.
- [2] Rawung JBM, Indrasti R, Bakrie B. Sustainable Agricultural Bioindustry Development: Integration of Cassava Cultivation with Beef Cattle Husbandry in North Sulawesi Province. *International Journal of Environment, Agriculture and Biotechnology*. 2018;3(4):264410.
- [3] Eduardo M, Svanberg U, Oliveira J, Ahrné L. Effect of Cassava Flour Characteristics on Properties of Cassava-Wheat-Maize Composite Bread Types. *International Journal of Food Science*. 2013;2013:305407.
- [4] Abass AB, Awoyale W, Sulyok M, Alamu EO. Occurrence of regulated mycotoxins and other microbial metabolites in dried cassava products from nigeria. *Toxins*. 2017;9(7):207.
- [5] Guo M, Song W, Buhain J. Bioenergy and biofuels: History, status, and perspective. *Renewable and sustainable energy reviews*. 2015;42:712-25.
- [6] Das I, Rakshit S. Millets, their importance, and production constraints. *Biotic stress resistance in millets*: Elsevier; 2016. p. 3-19.
- [7] Aworh OC. From lesser-known to super vegetables: the growing profile of African traditional leafy vegetables in promoting food security and wellness. *Journal of the Science of Food and Agriculture*. 2018;98(10):3609-13.
- [8] Ajayi, E.I.O.; Modo, E.U.; Kiakubu, O.T.; Molehin, O.R. Diabetes Care and Wound Healing Using *Nauclea latifolia* Manihot esculenta, and Other Natural Products. In *Bioactive Food as Dietary Interventions for Diabetes*; Academic Press: Cambridge, MA, USA, 2019. 545-58.
- [9] Ajayi EI, Molehin OR, Oloyede OI, Kumar V, Amara VR, Kaur J, et al. Liver mitochondrial membrane permeability modulation in insulin-resistant, uninephrectomised male rats by *Clerodendrum volubile* P. Beauv and *Manihot esculenta* Crantz. *Clinical Phytoscience*. 2019;5(1):1-9.
- [10] Mosayyebi B, Imani M, Mohammadi L, Akbarzadeh A, Zarghami N, Edalati M, et al. An update on the toxicity of cyanogenic glycosides bioactive compounds: Possible clinical application in targeted cancer therapy. *Materials Chemistry and Physics*. 2020;246:122841.
- [11] Nassar NMA, Graciano-Ribeiro D, Fernandes SDC and Araujo PC (2008). Anatomical alterations due to polyploidy in cassava, *Manihot esculenta* Crantz. *Genet. Mol. Res.* 7: 276-283. <http://dx.doi.org/10.4238/vol7-2gmr399>.
- [12] Graciano-Ribeiro D, Hashimoto-Freitas DY and Nassar NMA (2016). Comparative petiole anatomy of cassava (*Manihot*) species. *Genetics and Molecular Research* 15 (1): 1-13 DOI <http://dx.doi.org/10.4238/gmr.15017495>.
- [13] Fukuda, W.M.G., C.L. Guevara, R. Kawuki, and M.E. Ferguson. Selected morphological and agronomic descriptors for the characterization of cassava. *International Institute of Tropical Agriculture (IITA)*, Ibadan, Nigeria. 2010. 1-19.
- [14] Ndukwu B, Okoli B. Studies on Nigeria *Cucurbita moschata*. *Nigerian Journal of Botany*. 1992;5:18-26.
- [15] Pushpalatha R, Gangadharan B. Is cassava (*Manihot esculenta* Crantz) a climate “smart” crop? A review in the context of bridging future food demand gap. *Tropical Plant Biology*. 2020;13(3):201-11.
- [16] Ovalle TM, Vásquez-Ordóñez AA, Jimenez J, Parsa S, Cuellar WJ, Becerra Lopez-Lavalle LA. A simple PCR-based method for the rapid and accurate identification of spider mites (Tetranychidae) on cassava. *Scientific Reports*. 2020;10(1):1-10.
- [17] Zou Z, Yang J. Genomics analysis of the light-harvesting chlorophyll a/b-binding (Lhc) superfamily in cassava (*Manihot esculenta* Crantz). *Gene*. 2019;702:171-81.
- [18] Elsam JDE, del Carmen Peña-Chocarro M, Mereles F, Céspedes G. *Manihot takape* sp. nov. (Euphorbiaceae), a new tuberous subshrub from the Paraguayan Chaco. *PhytoKeys*. 2018(103):1.
- [19] Graciano-Ribeiro D, Hashimoto-Freitas D, Nassar N. Comparative petiole anatomy of cassava (*Manihot*) species. *Genetics and Molecular Research*. 2016;15:1-13.
- [20] Demotes-Mainard S, Péron T, Corot A, Bertheloot J, Le Gourrier J, Pelleschi-Travier S, et al. Plant responses to red and far-red lights, applications in horticulture. *Environmental and experimental Botany*. 2016;121:4-21.
- [21] Crang R, Lyons-Sobaski S, Wise R. *Epidermis*. *Plant anatomy*: Springer; 2018. p. 279-318.

- [22] Cach-Pérez MJ, Andrade JL, Cetzal-Ix W, Reyes-García C. Environmental influence on the inter-and intraspecific variation in the density and morphology of stomata and trichomes of epiphytic bromeliads of the Yucatan Peninsula. *Botanical Journal of the Linnean Society*. 2016;181(3):441-58.
- [23] Soengas P, Rodríguez VM, Velasco P, Cartea ME. Effect of temperature stress on antioxidant defenses in *Brassica oleracea*. *Acs Omega*. 2018;3(5):5237-43.
- [24] Asgarpour R, Ghorbani R, Khajeh-Hosseini M, Mohammadvand E, Chauhan BS. Germination of spotted spurge (*Chamaesyce maculata*) seeds in response to different environmental factors. *Weed Science*. 2015;63(2):502-10.
- [25] Toscano S, Ferrante A, Romano D. Response of Mediterranean ornamental plants to drought stress. *Horticulturae*. 2019;5(1):6.
- [26] Derso C, Mahmud A. Study on morphological characters of four cassava (*Manihot esculenta* Crantz) varieties as cultivated in Fafen District, Ethiopian Somali Regional State. *Asian J of Biotechnology and Bioresource Technology* doi. 2018;10.
- [27] Neto ILdC, Martins FM, Martins MLL, de Oliveira Silva LR. Comparative leaf anatomy of wild *Manihot* Mill. species (Euphorbiaceae) from Chapada Diamantina, Bahia, Brazil. *Nordic Journal of Botany*. 2017;35(2):207-19.
- [28] Santos AS, Martins MLL, Martins FM, de Carvalho PC, da Silva Ledo CA, Amorim AM. A new species and putative hybrids of *Manihot* (Euphorbiaceae) from Brazil. *Plant Systematics and Evolution*. 2019;305(8):663-74.
- [29] Silva LRdO, Martins FM, Martins MLL. Anatomy and histochemistry of *Manihot caerulescens* Pohl (Euphorbiaceae) Leaf anatomy of *Manihot caerulescens*. *Feddes Repertorium*. 2019;130(4):379-88.